

Perturbative moduli stabilisation in type IIB/F-theory

Yifan Chen

LPTHE, Sorbonne Université

arXiv: 1803.08941,

Ignatios Antoniadis, Y. C, George K. Leontaris.

String Phenomenology 2018

Warsaw

04.07.2018

Outlines

- 1 Moduli stabilisation
- 2 String Loop Correction
- 3 Single $D7$ -brane
- 4 Three Intersecting $D7$ -branes
- 5 D-term
- 6 Summary
- 7 Backup

Moduli Fields

- In string theory, all **scales, particle spectra and couplings** come from the **geometry of the extra dimensions**.
- The geometry of the extra dimensions is parameterised by complex scalar fields: Moduli fields.
- Massless moduli generate **long-range fifth forces** and lead to theory **outside the regime of validity**.
- Moduli stabilisation: stabilise the geometry of the extra dimensions which leads to a vacuum that can embed Standard Model matter spectrum in four dimensions.

Type IIB Moduli Stabilisation

In IIB string theory, the moduli definitions are

- Axion-dilaton (**String coupling**): $S = e^{-\phi} + iC_0 \equiv 1/g_s + iC_0$,
- Complex structure (**Shape**) Moduli: z_a ,
- Kähler (**Size**) Moduli: complexified 4-cycle volumes T_i .

One can turn on the 3-form fluxes in the superpotential
[Gukov-Vafa-Witten '99]:

$$\mathcal{W}_0 = \int G_3 \wedge \Omega,$$

where $G_3 = F_3 - SH_3$, Ω contains z_a .

- **Supersymmetric condition** fixes the first two:

$$D_S \mathcal{W}_0 = D_{z_a} \mathcal{W}_0 = 0.$$

No Scale-Structure

After fixing all the complex moduli and axion-dilaton, only Kähler moduli left... Tree level Kähler potential is:

$$\mathcal{K}_0(T_i) = - \sum_{i=1}^3 \ln(-i(T_i - \bar{T}_i))$$

which leads to:

$$V = e^{\mathcal{K}} \left(\sum_{I, J \neq T_i} D_I \mathcal{W} \mathcal{K}_{I\bar{J}}^{-1} D_{\bar{J}} \mathcal{W} + \sum_{T_i} D_{T_i} \mathcal{W} \mathcal{K}_{T_i \bar{T}_i}^{-1} D_{\bar{T}_i} \mathcal{W} - 3|\mathcal{W}|^2 \right) = 0.$$

- Vanishing vacuum energy with T_i unstabilised,
- Broken SUSY in T_i .

Breaking No Scale-Structure

- KKLT: **Non-perturbative effects** [Kachru-Kalosh-Linde-Trivedi '03]

$$\mathcal{W} = \mathcal{W}_0 + \sum_{i=1}^{h_+^{1,1}} \Lambda_i e^{-\lambda_i T_i} .$$

Supersymmetric AdS minimum, uplifted by $\overline{D3}$ -branes.

- LVS: **Non-perturbative effects** + α' corrections
[Balasubramanian-Berglund-Conlon-Quevedo '05]

$$\begin{aligned} \mathcal{K}_{LVS} &= -2 \ln(\tau_b^{\frac{3}{2}} - \tau_s^{\frac{3}{2}} + \xi), \\ \mathcal{W}_{LVS} &= \mathcal{W}_0 + \Lambda e^{-\lambda \tau_s}. \end{aligned}$$

F-term spontaneous supersymmetry breaking, uplifted by D-term.

Perturbative Corrections

α' corrections $\hat{\xi}$ and string one-loop corrections $\hat{\delta}$ arise both in the string frame as corrections to the Einstein kinetic terms:

$$\left[e^{-2\phi}(\mathcal{V} + \hat{\xi}) + \hat{\delta} \right] \mathcal{R},$$

which can be accounted by the shifts:

$$\mathcal{V} \rightarrow \mathcal{V} + \hat{\xi} \quad ; \quad e^{-2\phi_4} = e^{-2\phi} \mathcal{V} \rightarrow e^{-2\phi_4} + \hat{\delta}.$$

The radiative corrected Kähler potential reads:

$$\begin{aligned} \mathcal{K} &= -2 \ln \left[e^{-2\phi}(\mathcal{V} + \hat{\xi}) + \hat{\delta} \right] \\ &= -\ln(S - \bar{S}) - 2 \ln \left(\hat{\mathcal{V}} + \xi + \delta \right), \end{aligned}$$

String One-Loop Corrections with $D7$ -Branes

- In IIB string theory, graviton vertices from the 4d Einstein action localised in the internal space emit closed strings in the bulk carrying **transverse momentum p_\perp** .
- p_\perp is not conserved and can flow on the other boundary because there are **local tadpoles (point-like sources in the transverse dimensions)**. These contribute to the amplitude in the large transverse volume V_\perp limit: **[Antoniadis-Bachas '98]**

$$\mathcal{T}(\mathcal{A}) \sim \frac{1}{V_\perp} \sum_{|p_\perp| < M_s} \frac{1}{p_\perp^2} F(\vec{p}_\perp),$$

where the $F(\vec{p}_\perp)$ are the local tadpoles in the momentum space and

$$\vec{p}_\perp = \left(\frac{n_1}{R}, \dots, \frac{n_d}{R} \right)$$

- **2D summation** leads to **logarithmic divergence**.

A Single D7-Brane

Single D7-brane

- Kähler modulus: the world volume part τ and the transverse part u :

$$\mathcal{K} = -2 \ln(\tau\sqrt{u} + \xi + \eta\ln(u)).$$

- Perturbative region:

$$|\eta\ln(u)| \ll \tau\sqrt{u}.$$

- Large volume expansion of the derivative with respect to u :

$$\frac{dV_F(\tau, u)}{du} = -\eta\mathcal{W}_0^2 \frac{3(-10 + 3\ln(u))}{4\tau^3 u^{5/2}} + O(\eta^2) + O(\xi).$$

- For $\eta < 0$, the transverse direction u can have a minimum, while world volume part τ not.

Three Intersecting $D7$ -Branes

Three Intersecting D7-Branes

- Now the Kähler potential reads:

$$\begin{aligned}\mathcal{K} &= -2\ln(\sqrt{\tau_1\tau_2\tau_3} + \sum_i 2\eta_i \ln(\frac{\mathcal{V}}{\tau_i})) \\ &= -2\ln(\sqrt{\tau_1\tau_2\tau_3} + \sum_i \eta'_i \ln(\tau_i)), \quad \eta'_a = \sum_i \eta_i - 2\eta_a.\end{aligned}$$

- The derivative with respect to either τ_a :

$$\frac{dV_F(\tau_i)}{d\tau_a} = \mathcal{W}_0^2 \frac{3(\sum_{i \neq a} (8\eta'_i - 3\eta'_i \ln(\tau_i)) + 10\eta'_a - 3\eta'_a \ln(\tau_a))}{4 \prod_{i \neq a} \tau_i^{\frac{3}{2}} \tau_a^{\frac{5}{2}}} + O(\eta'^2).$$


- Minimisation condition:

$$\eta_1 = \eta_2 = \eta_3 = \eta_\tau < 0,$$

- But the minimum is only for the total volume, since:

$$\mathcal{K} = -2\ln(\mathcal{V} + 2\eta_\tau \ln(\mathcal{V})).$$

$$V_F(\mathcal{V}) = \frac{\eta_\tau \mathcal{W}_0^2}{\mathcal{V}^3} (3\ln(\mathcal{V}) - 12) + \mathcal{O}(\eta_\tau^2)$$



The minimum is inside the **perturbative region** and SUSY is **spontaneously broken!**

D-term from $D7$ -Branes

- There remain two moduli, the ratios, unfixed.
- A deviation from the condition $\eta_1 = \eta_2 = \eta_3$ will destabilise the vacuum.
- Solution: Introduce world-volume dependent D-term from each $D7$ -brane for both uplifting and ratios stabilisation.

$$V_{D_a} = \frac{d_a}{\tau_a} \left(\frac{\partial K}{\partial \tau_a} \right)^2 = \frac{d_a}{\tau_a^3} + O(\eta_i),$$

- The sum of the potential leads to a GUT scale compactification volume and stabilise the ratios:

$$\tau_a = \left(\frac{d_a^3}{d_1 d_2 d_3} \right)^{\frac{1}{9}} \mathcal{V}_0^{\frac{2}{3}} \quad \ln(\mathcal{V}_0) \simeq 5.$$

$$\frac{M_{pl}}{\mathcal{V}_0} \simeq \Lambda_{GUT}.$$

Conclusion

- The string loop corrections display **logarithmic dependence on the size of the moduli transverse to $D7$ -branes** in the presence of **local tadpoles**. With 3 intersecting $D7$ -branes, one can **stabilise the total internal volume**.
- Magnetised $D7$ -branes, which can generate D-terms that **depend on the world-volume** of the corresponding $D7$ -branes, can **stabilise the ratios and uplift the potential to de-Sitter minima** where both F-term and D-term supersymmetry are spontaneously broken.
- The realisation of this geometric stabilisation mechanism and the uplifting is a viable scenario in F-theory. Thus one can **use the F-theory model building to realise the Standard Model**.

*Thank
you!*

Backup

Three Intersecting $D7$ -Branes

- The whole 6-dimensional volume as triple products of 2-cycle moduli:

$$\mathcal{V} = \frac{1}{6} \kappa_{abc} v^a v^b v^c,$$

- In the framework of 3 intersecting $D7$ -branes, take 2-cycle v^a as the transverse volume modulus of each $D7$ -brane with world-volume τ_a :

$$v_a = \frac{\mathcal{V}}{\tau_a},$$

- Take κ_{abc} as ϵ_{abc} for simplicity. Then the volume can be expressed as

$$\mathcal{V} = v_1 v_2 v_3 = \sqrt{\tau_1 \tau_2 \tau_3}$$